

Reconceptualizing the hyporheic zone for nonperennial rivers and streams

Amanda G. DelVecchia^{1,16}, Margaret Shanafield^{2,17}, Margaret A. Zimmer^{3,18}, Michelle H. Busch^{4,19}, Corey A. Krabbenhoft^{5,20}, Rachel Stubbington^{6,21}, Kendra E. Kaiser^{7,22}, Ryan M. Burrows^{8,23}, Jake Hosen^{9,24}, Thibault Datry^{10,25}, Stephanie K. Kampf^{11,26}, Samuel C. Zipper^{12,27}, Ken Fritz^{13,28}, Katie Costigan^{14,29}, and Daniel C. Allen^{15,30}

¹Department of Biology, Duke University, 130 Science Drive, Durham, North Carolina 27708 USA

²College of Science and Engineering, Flinders University, Ring Road, Bedford Park, South Australia 5042 Australia

³Department of Earth and Planetary Sciences, 1156 High Street, University of California, Santa Cruz, California 95064 USA

⁴Department of Biology, University of Oklahoma, 730 Van Vleet Oval, Norman, Oklahoma 73019 USA

⁵Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, 2003 Upper Buford Circle, St Paul, Minnesota 55108 USA

⁶School of Science and Technology, Nottingham Trent University, Clifton Campus, Clifton Lane, Nottingham NG11 8NS United Kingdom

⁷Geosciences Department, Boise State University, 1295 University Drive, Boise, Idaho 83725 USA

⁸School of Ecosystem and Forest Sciences, University of Melbourne, 500 Yarra Boulevard, Burnley, Victoria 3121 Australia

⁹Department of Forestry and Natural Resources, Purdue University, 195 Marsteller Street, West Lafayette, Indiana 47906 USA

¹⁰French National Institute for Agriculture, Food, and the Environment, UR-RiverLy, Centre de Lyon-Villeurbanne, Centre Lyon-Grenoble Auvergne-Rhône-Alpes, 5 rue de la Doua CS70077, 69626 Villeurbanne CEDEX France

¹¹Department of Ecosystem Science and Sustainability, Colorado State University, 1476 Campus Delivery, Fort Collins, Colorado 80521 USA

¹²Kansas Geological Survey, 1930 Constant Avenue, Lawrence, Kansas 66047 USA

¹³Office of Research and Development, United States Environmental Protection Agency, 26 West Martin Luther King Drive, Mailstop 585, Cincinnati, Ohio 45268 USA

¹⁴School of Geosciences, University of Louisiana, 611 McKinley Street, Hamilton Hall 323, P.O. Box 43717, Lafayette, Louisiana 70504 USA

¹⁵Department of Ecosystem Science and Management, The Pennsylvania State University, 311 Forestry Resources Building, University Park, Pennsylvania 16802 USA

Abstract: Nonperennial streams dominate global river networks and are increasing in occurrence across space and time. When surface flow ceases or the surface water dries, flow or moisture can be retained in the subsurface sediments of the hyporheic zone, supporting aquatic communities and ecosystem processes. However, hydrological and ecological definitions of the hyporheic zone have been developed in perennial rivers and emphasize the mixing of water and organisms from both the surface stream and groundwater. The adaptation of such definitions to include both humid and dry unsaturated conditions could promote characterization of how hydrological and biogeochemical variability shape ecological communities within nonperennial hyporheic zones, advancing our understanding of both ecosystem structure and function in these habitats. To conceptualize hyporheic zones for nonperennial streams, we review how water sources and surface and subsurface structure influence hydrological and physicochemical conditions. We consider the extent of this zone and how biogeochemistry and ecology might

**This section of the journal is for the expression of new ideas, points of view, and comments on topics of interest to aquatic scientists. The editorial board invites new and original papers as well as comments on items already published in Freshwater Science. Format and style may be less formal than conventional research papers; massive data sets are not appropriate. Speculation is welcome if it is likely to stimulate worthwhile discussion. Alternative points of view should be instructive rather than merely contradictory or argumentative. All submissions will receive the usual reviews and editorial assessments.*

E-mail addresses: ¹⁶amanda.delvecchia@duke.edu; ¹⁷margaret.shanafield@flinders.edu.au; ¹⁸margaret.zimmer@ucsc.edu; ¹⁹buschmh@ou.edu; ²⁰cakrabbe@gmail.com; ²¹rachel.stubbington@ntu.ac.uk; ²²kendrakaiser@boisestate.edu; ²³ryan.burrows@unimelb.edu.au; ²⁴hosen@purdue.edu; ²⁵thibault.datry@inrae.fr; ²⁶stephanie.kampf@colostate.edu; ²⁷samzipper@ku.edu; ²⁸fritz.ken@epa.gov; ²⁹costigan@louisiana.edu; ³⁰daniel.c.allen@psu.edu

Received 17 May 2021; Accepted 14 December 2021; Published online 22 April 2022. Associate Editor, Matthew Baker.

Freshwater Science, volume 41, number 2, June 2022. © 2022 The Society for Freshwater Science. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0), which permits non-commercial reuse of the work with attribution. For commercial use, contact journalpermissions@press.uchicago.edu. Published by The University of Chicago Press for the Society for Freshwater Science. <https://doi.org/10.1086/720071>

vary with surface states. We then link these components to the composition of nonperennial stream communities. Next, we examine literature to identify priorities for hydrological and ecological research exploring nonperennial hyporheic zones. Lastly, by integrating hydrology, biogeochemistry, and ecology, we recommend a multidisciplinary conceptualization of the nonperennial hyporheic zone as the porous subsurface streambed sediments that shift between lotic, lentic, humid, and dry conditions in space and time to support aquatic–terrestrial biodiversity. As river drying increases in extent because of global change, we call for holistic, interdisciplinary research across the terrestrial and aquatic sciences to apply this conceptualization to characterize hyporheic zone structure and function across the full spectrum of hydrological states.

Key words: aquatic–terrestrial transition zone, dry rivers, ecotone, hyporheic, intermittent river, intermittent stream, nonperennial stream, ephemeral stream, riverbed sediments, subsurface sediments, temporary river, temporary stream

Nonperennial streams and rivers are dynamic ecosystems in which surface flow ceases and most or all surface water is lost at some point in space and time (Datry et al. 2017; Fig. 1A–C). Nonperennial streams dominate global river networks (Messenger et al. 2021), and both their occurrence and extent are increasing because of climate change, land-use change, and human demand for freshwater (Datry et al. 2014, Perkin et al. 2017, Allen et al. 2019). Natural stream drying generates and maintains habitat heterogeneity and extensive aquatic–terrestrial linkages that promote biodiversity and biogeochemical complexity (Datry et al. 2014), providing ecosystem goods and services (Acuña et al. 2014, Datry et al. 2018, Stubbington et al. 2020). Nonperennial stream reaches that are dry at the surface might retain subsurface water as vapor or liquid, the latter sometimes flowing. Thus, the subsurface environment can support abundant and diverse aquatic communities (Williams 1996, Febria et al. 2015) and ecological functions beneath both wet and dry channels (Burrows et al. 2017, Colls et al. 2019).

The hyporheic zone (HZ) allows hydrological and ecological connectivity to exist between surface and subsurface environments. The HZ term originated when Orghidan (1959) examined the fauna within a hole in a streambed,

noting that the subsurface sediments provided an ecotone that mixed the physical (temperature, water velocity, light) and chemical (organic content, dissolved oxygen [DO], salinity) conditions of waters originating from both the surface and the groundwater. Importantly, Orghidan (1959) also recognized this zone as a transition from aquatic (saturated) to terrestrial (unsaturated) hydrological conditions. As the number of HZ-focused studies has grown over the past decade (Krause et al. 2011, Ward 2016, Woessner 2017), a range of hydrological and ecological conceptualizations and definitions have been developed for perennial streams (Krause et al. 2009, Cardenas 2015). Hydrological conceptualizations of the HZ typically refer to the mixing of surface water and groundwater and (usually implicitly) assume saturated conditions in the streambed (Gooseff et al. 2003, Runkel et al. 2003, Stonedahl et al. 2010, Boano et al. 2014). Similarly, biological definitions emphasize inhabitation by aquatic communities, including organisms that inhabit the surface stream and groundwater (Stanford et al. 1994, Boulton 2000). However, HZs in nonperennial streams can violate these restrictive conceptualizations because of their variable—and sometimes nonexistent—contributions of surface water and groundwater, resultant range of saturated to



Figure 1. Instream states vary between and within seasons in a winterbourne chalk stream in England, United Kingdom: a winter flowing phase (A), a summer low-flow phase (B), and a summer dry phase (C). Photographs supplied by the United Kingdom Environment Agency under the Open Government License version 3.0.

dry conditions, and corresponding range of resident taxa and biogeochemical conditions. Nonperennial stream HZs, thereby, encompass aquatic and terrestrial conditions, limiting applicability of methods and relevant metrics from either science across the full hydrograph. This cross-disciplinary narrow focus on saturated states and aquatic organisms, as well as independent and potentially divergent discipline-specific definitions of the HZ, fail to represent the structure and function of nonperennial HZs. Thus, interdisciplinary understanding of the biogeochemical, ecological, and hydrological roles of the HZ has been limited.

Here, we compare and synthesize existing HZ definitions and case studies from hydrological, biogeochemical, and ecological research in nonperennial streams to develop a more inclusive conceptualization of the HZ that focuses on its distinct role as dynamic subsurface habitat. First, we investigate how water sources, flowpath directions, and residence times drive the state, extent, and saturation conditions of the HZ (Fig. 2). Second, we synthesize how physicochemical conditions and microbially mediated biogeochemical processes may vary with the degree of saturation, water source, and flowpath length, thereby affecting habitat conditions as well as chemical and biological interactions with the surface stream. Third, we explore the effects of hydrological, physicochemical, and biogeochemical conditions on invertebrate and vertebrate communities and biodiversity in nonperennial HZs. We then conduct a literature review to identify gaps representing priorities for interdisciplinary nonperennial HZ research and propose methods to consistently investigate relationships between hydro-

logy, biogeochemistry, and ecology in these widespread, understudied systems. Integrating insights from across disciplines, we propose a unified definition of the nonperennial HZ that explicitly recognizes these subsurface sediments as dynamic ecotones shifting between flowing, wet, and dry states that support aquatic–terrestrial biodiversity and, thus, ecosystem functioning.

HOW DO SURFACE WATER AND GROUNDWATER HYDROLOGY AFFECT THE STATE, EXTENT, AND SATURATION CONDITIONS OF THE HZ?

In both perennial and nonperennial streams, water movement into, through, and out of the HZ is controlled by the geomorphic and hydraulic properties of the streambed, its wider corridor (Ward and Packman 2019), and the hydraulic gradient between the stream and underlying groundwater (Winter 1999). Pressure gradients force surface water into and through the subsurface, where flowpaths are controlled by the geomorphic and hydrological characteristics of the stream (Boano et al. 2014, Zimmer and Lautz 2014). Hydrological studies of the HZ often focus on short (0.1–1 m) subsurface flowpaths caused by these pressure-induced interactions (Lewandowski et al. 2019). Depending on research objectives, hydrologists typically define the HZ based on surface water–groundwater interactions (e.g., Woessner 2017), transport of solutes, including nutrients (Triska et al. 1989, Bencala et al. 2011), or solely on surface water circulation (Ward 2016). The recent synthesis of Lewandowski et al. (2019) summarizes the breadth of current




Figure 2. The hyporheic zone, or porous sediments beneath and surrounding the streambed, can contain contributions from various water sources (e.g., surface water, regional groundwater). All flowing water moves along flowpaths. Although depicted in blue, the hyporheic zone is not always saturated.

hydrological definitions by considering the HZ as a zone of saturated, porous streambed sediments in which at least 10% surface water and groundwater interact, or in which flowpaths begin and return to the surface. However, no hydrological definition of the HZ accommodates the dynamic, spatially and temporally discontinuous flow regimes of nonperennial streams.

In contrast to perennial stream HZs, the water content, or 3-dimensional spatial extent of water within interstitial spaces (i.e., saturation), can change profoundly during the hydrological cycle of a nonperennial stream. Nonperennial streams with connection between surface water and groundwater might disconnect seasonally, the HZ potentially comprising a 3-dimensional patchwork of both saturated and unsaturated sediments during dry seasons (Fig. 3E, F) (Fleckenstein et al. 2006). Perched aquifers will likely fill quickly and saturate during periods in which the stream is flowing but slowly desaturate because of infiltration and evapotranspiration when the surface streambed is dry (Villeneuve et al. 2015; Fig. 3C, D). Stream reaches with bypass (or parafluvial) flow, defined as water that circumvents channels via subsurface pathways, would likely have spatially variable saturation due to preferential pathways and surface states (Fig. 3A, B).

The lateral vs vertical extent of the HZ varies considerably, depending on geomorphic landscape characteristics (Stanford and Ward 1993). In gaining reaches (e.g., due to seasonally high groundwater levels or where a perched water table or other geological feature causes persistent inflow), shallow water tables may be more laterally extensive (Fleckenstein et al. 2006). Where floodwaters have overtopped stream banks, return flows to the stream might also occur for relatively long periods (McCallum and Shanafield 2016). Conversely, the HZ is laterally narrow when streambed hydraulic conductivity is high, floodplains are narrow or constricted by bedrock, there is strong groundwater inflow, or a vertically disconnected water table drives a strong losing hydraulic gradient (Wondzell and Swanson 1999, Kiel and Cardenas 2014).

As nonperennial streams shift between 3 surface states: flowing, ponded, and dry (Shanafield et al. 2021), hydrological, biogeochemical, and biological fluxes within the HZ vary depending on stream geology and geomorphology and the direction of hydrological exchange. When surface flow ceases, connected or isolated surface water pools and extensive dry conditions may occur in the surface channel, with concurrent changes in the magnitude, or even occurrence, of subsurface water fluxes. In gaining reaches, the contribution of groundwater can keep the HZ saturated and maintain hyporheic flowpaths during periods without surface flow or water, potentially resulting in persistent isolated pools. In contrast, in a losing stream, the streambed sediments can quickly become unsaturated because of the lack of connection between the stream and groundwater




Figure 3. Three exemplary scenarios in which the hyporheic zone (HZ) of nonperennial streams deviates from traditional conceptualizations. In the parafluvial/bypass flow scenario, only a fraction of the HZ might comprise saturated sediments, which may provide refuge for hyporheic communities when the stream ceases to flow (A, B). Where groundwater tables are shallow, the subsurface may remain humid all year. In the perched scenario, surface water inputs to the HZ might be nonexistent after the stream stops flowing or dries, but interstitial water content could be sustained by water retained above the confining layer (represented by brown) (C, D). In the losing disconnected scenario, disconnected pools might be connected by hyporheic exchange (E, F).

(Falke et al. 2011; Fig. 3E, F), although shallow groundwater can influence surface and shallow subsurface flow even when the water table is not hydraulically connected to the stream (Quichimbo et al. 2020; Fig. 3C, D). Where bypass flow occurs, surface water may enter the streambed at an upstream location, exiting and reactivating surface flow at some downstream point, even if the surface sediments in between are dry (Fig. 3A, B) (Costigan et al. 2015, Busch et al. 2020). In each of these situations, water physicochemistry depends on mixing between surface water and groundwater, but literature on hyporheic hydrology rarely defines the term groundwater.

Different groundwater sources to hyporheic sediments have varying impacts on biogeochemical cycling. Regional groundwater that has been in long-term contact with subsurface geology will reflect the characteristic geochemistry of the parent material (Appelo and Postma 2004). In contrast, groundwater flowing through short within-catchment flowpaths (e.g., from subsurface flow during storms or snowmelt) may be compositionally comparable to surface water. For example, springs and seeps drive gaining conditions in some nonperennial stream systems, contributing water that reflects aquifer chemistry (Wroblicky et al. 1998). These differences in the source and age of water entering and exiting nonperennial stream channels influence hyporheic water physicochemistry, including temperature and solutes and, therefore, habitat for biota and potential for nutrient processing within the HZ (Brookfield et al. 2021). Differences in groundwater sources could be especially influential in nonperennial HZs, where surface water contributions could be completely absent at times.

Thus, hydrological conditions in the HZ are particularly dynamic in nonperennial streams, in which drivers of water presence or absence are complex in both time and space (Hammond et al. 2021). However, such variability is at odds with existing definitions of the HZ. If a surface stream is not flowing or is dry (with or without pools), classical hydrological definitions of surface–subsurface exchange flows are not met, but the subsurface sediments may remain hydrologically active. The requirement for the HZ to be saturated, by definition, thus excludes nonperennial HZs that comprise saturated or unsaturated subsurface sediments.

HOW DO SPATIAL AND TEMPORAL LINKAGES BETWEEN SURFACE WATER AND GROUNDWATER AFFECT BIOGEOCHEMISTRY AND MICROBIAL ECOLOGY?

Hydrology affects HZ biogeochemistry by controlling both water sources (surface water, shallow and deep groundwater) and how water travels through the sediment matrix and, thus, interacts with microbial communities. Just as conditions in nonperennial surface streams occur along a continuum encompassing flowing, lentic, and dry states, the HZ

ranges between saturated (flowing or lentic), unsaturated, and dry conditions. In saturated flowing conditions, dominant microbial processes change along subsurface flowpaths (Febria et al. 2012, Mach et al. 2015). Transitions between states and source inputs influence sediment moisture, temperature, DO concentrations, redox conditions, specific conductance, and pH (Gómez-Gener et al. 2016). These physicochemical conditions shape and are shaped by microbial communities and associated biogeochemical processes (Sabater et al. 2016), thereby providing a mechanism by which hyporheic hydrology regulates the biogeochemistry of nonperennial streams.

Different water sources can affect physicochemical conditions and biogeochemical processes in the HZ. In some settings, upwelling groundwater can be more NO_3^- rich, P rich, and thermally stable than surface water (Boulton et al. 2010) and can reduce interstitial sedimentation. In contrast, downwelling surface water is often more oxygenated and richer in organic matter than groundwater (Boulton et al. 2010). Interactions between these different water sources can make the HZ an ecotone between oxidizing and reducing conditions (Krause et al. 2011). The characteristics of water from different sources also vary with flow conditions. Downwelling water can carry and deposit fine sediment and biomass that clogs interstices, especially during high-flow conditions, reducing hydraulic conductivity and creating hyporheic conditions favorable to anoxia (Brunke and Gonser 1997). During low surface flows, dense growth of heterotrophic and autotrophic biofilms can cover or infiltrate the HZ, also reducing hydrologic exchange (Stanley et al. 1997, Caruso et al. 2017).

Both the magnitude of hydrologic exchange and conditions in drying surface water can influence conditions in the HZ. As nonperennial streams transition from flowing to lentic, environmental conditions in any persisting surface pools can diverge after hydrological connectivity is lost (Verdonschot et al. 2015, Casas-Ruiz et al. 2016, Fig. 4A, B). High water-residence times in these pools, combined with high temperatures, can fuel gross primary production (GPP) and ecosystem respiration (ER). If light availability is high, GPP:ER ratios can also be high (Casas-Ruiz et al. 2016), and, thus, any water infiltrating the HZ will have higher DO concentrations. In contrast, high dissolved organic C concentrations, such as from leaf leachates and soils (Chafiq et al. 1999, Siebers et al. 2016), light limitation (Casas-Ruiz et al. 2016, Hensley et al. 2019, Hosen et al. 2019), or high decomposer activity (Abril et al. 2016), can cause low GPP:ER ratios in pools as heterotrophic processes (e.g., ammonification; Skoulikidis et al. 2017) dominate and gradually reduce DO concentrations (Stanley et al. 1997, von Schiller et al. 2011).

In both perennial and nonperennial streams, inputs of DO, organic matter, and other terminal electron acceptors (e.g., NO_3^- and SO_4^{2-}) act within the physical structure




Figure 4. Along flowpaths, the concentrations of various terminal electron acceptors used in respiration are depleted as concentrations of reduced byproducts increase (modified from Kehew 2001 and Schlesinger and Bernhardt 2013), as might occur along a subsurface flowpath from the upstream to downstream piezometers in either connected (A) or disconnected (B) streams. In panel A, surface water is clearly connected, as are subsurface flowpaths. When surface water is disconnected (B), subsurface flowpaths might persist with these concomitant changes in constituents. However, the concentrations of constituents entering the hyporheic zone might differ (blue during the most connected periods, ranging to red as the stream begins to dry; C). Infiltrating water from disconnected pools might have higher NH_4^+ concentrations, more variation in dissolved oxygen concentrations, higher dissolved organic matter, and higher NO_3^- , all of which is depleted at varying rates over time in the hyporheic zone. As the hyporheic zone becomes disconnected from surface water, it is possible that water chemistry will more closely resemble water with a longer residence time.

provided by sediment, affecting interstitial physicochemistry, microbial activity, and biogeochemical processes along hyporheic flowpaths (Gilbert et al. 1990, Vervier et al. 1992, Hancock et al. 2005, Claret and Boulton 2008, Zlatanović et al. 2018, Fig. 4C). Nonperennial streams can have especially variable subsurface flowpath lengths and inputs of organic matter and nutrients. For example, in nonperennial streams with bypass flow, water penetrates further into the subsurface as surface flow increases, creating longer flowpaths (Vázquez et al. 2007). Longer flowpaths in losing reaches can have higher mineralization of dissolved organic matter, leading to more reduced but bioavailable forms of inorganic N (i.e., NH_4^+), or potentially storing infiltrated C and N as microbial biomass and dissolved gases (Newcomer et al. 2018). As demonstrated in gravel-bed floodplains, long flowpaths that contribute to nonperennial spring flow can have extremely low dissolved organic C concentrations, low DO concentrations, chemoautotrophy (e.g., hydrogenotrophic methanogenesis, nitrification), and a tendency toward reducing reactions (e.g., Fe reduction, SO_4^{2-} reduction, and meth-

anogenesis) (Helton et al. 2015, DelVecchia et al. 2016). When these substrates encounter oxidizing conditions caused by surface water, they can enable heterotrophic and chemotrophic processes (Datry and Larned 2008).

As the surface channel dries, photosynthesis and substrate delivery to the HZ diminishes (Colls et al. 2019), altering the quality and quantity of C inputs to underlying HZs. Hyporheic heterotrophs, thus, tend to shift to less labile C sources (Granados et al. 2020). As the HZ itself dries, reduced substrate diffusion and microbial motility can directly inhibit microbial activity (Humphries and Baldwin 2003). However, if subsurface interstices retain moisture, organic matter decomposition and mineralization can continue (Solagaistua et al. 2016, von Schiller et al. 2017) and can even exceed surface rates if conditions are more favorable for biotic processes (Burrows et al. 2017, Arias-Real et al. 2020).

The changes in biogeochemical processes that occur along flowpaths are also mediated by moisture when water is not flowing. For example, N cycling varies depending on

whether 1) oxygenated humid conditions facilitate net nitrification, 2) deoxygenated humid conditions limit ammonification and nitrification but favor denitrification, or 3) dry conditions limit each of these processes (Tzoraki et al. 2007, Gómez et al. 2012, Sabater et al. 2016). Drying also has strong but unpredictable effects on sediment affinity for P. P mobility typically increases as redox conditions become more negative (Ann et al. 1999), but the presence of metals, such as Fe and Al, can modulate this process (Peng et al. 2007). As a result, stream drying can either increase or reduce sediment binding (Sabater et al. 2016, von Schiller et al. 2017). Thus, depending on local conditions, drying and rewetting processes can have contrasting outcomes for microbial communities and biogeochemical processes.

When the HZ retains some moisture, it can harbor desiccation-tolerant microbial taxa (Gionchetta et al. 2019). However, drying and rewetting also create strong selective pressures that eliminate some taxa (Zeglin et al. 2011, Febria et al. 2012, Timoner et al. 2014). The duration and severity of dry phases are master variables controlling non-perennial stream microbial communities (Sabater et al. 2016, Colls et al. 2019). If the HZ retains a connection to groundwater during dry periods, hyporheic sediments can remain saturated or humid, enabling the survival of many microbial taxa (Gionchetta et al. 2019). Humidity, redox conditions, and microbial habitat are controlled not only by water sources but by organic matter content and sediment structure, which are, thus, also key determinants of microbial community composition (Zlatanović et al. 2018, Arias-Real et al. 2020).

The hydrological and biogeochemical controls on microbial communities in non-perennial HZs are relatively well established, but the relative importance of different factors and how they interact to control community composition is less clear. Specifically, it remains unclear whether local environmental sorting or dispersal and biotic interactions (e.g., competition) drive community composition. Benthic community composition in non-perennial streams is primarily controlled by redox conditions (Gionchetta et al. 2020), and both the community composition and habitat conditions present before an HZ dries can influence respiration rates that affect dry-phase conditions and communities (Duarte et al. 2017, Newcomer et al. 2018), exemplifying the potential for environmental sorting. Drying alters dispersal patterns by changing the water sources supplying the HZ (Fazi et al. 2013). As a dry phase progresses, surface water inputs decrease, whereas deeper groundwater inputs can remain steady, decrease, or increase, changing the number and taxonomic composition of microbes that passively disperse to hyporheic sediments (Saup et al. 2019). Hydrological conditions likely modulate whether sorting or dispersal mechanisms dominate, with dispersal having more influence during high-flow conditions and environmental sorting and biotic interactions becoming more

important as water-residence times increase in drier conditions (Datry et al. 2016).

The biogeochemical transformations that occur during dry phases are likely to affect community recovery following subsequent rewetting. Microbial communities that survive in humid conditions in the HZ can rapidly recover in the surface when streams rewet (Timoner et al. 2020). This recovery can be enhanced by nutrients liberated from microbial cells that lysed during dry phases because of environmental stress (Birch and Friend 1956, Baldwin and Mitchell 2000, Leung et al. 2020). Nutrient mineralization in the HZ can thereby contribute to GPP and ER in the surface sediments of gaining reaches during rewetting events that re-establish vertical hydrological connectivity (Sabater et al. 2016). In addition, both discontinuous flow and long flowpaths create reducing subsurface environments, so flow resumptions that drive hyporheic contributions to the well-oxygenated surface can create potential control points for autotrophic production (Burrows et al. 2020), heterotrophy, chemotrophy, and microbial dispersal (Bernhardt et al. 2017, von Schiller et al. 2019). The contribution of reduced hyporheic substrates to surface water could affect ecological and biogeochemical processes that affect energy sources, contribute to overall ecosystem function, and potentially alter the quality of downstream and laterally connected waters (Brookfield et al. 2021).

HOW DO HYDROLOGY AND BIOGEOCHEMISTRY AFFECT THE ROLE OF THE HZ IN MAINTAINING BIODIVERSITY IN NONPERENNIAL STREAMS?

Biological definitions of the HZ typically describe the organisms that inhabit it based primarily on our understanding of aquatic invertebrate assemblages. Some such definitions are broad, with Stanford and Ward (1988) considering the HZ as those sediments “penetrated by riverine animals” (p. 64) and Lewandowski et al. (2019) recognizing the subsurface sediments as supporting a “characteristic hyporheic community” (p. 3). Others emphasize the HZ as inhabited by both temporary residents, which mainly live in either the benthic zone or groundwater, and permanent hyporheic specialists (Williams and Hynes 1974, Boulton 2000). All such ecological conceptualizations mirror hydrological definitions in both their ecotonal aspects (i.e., mixing of water or organisms from the surface stream and groundwater; Brunke and Gonser 1999) and their explicit or implicit consideration of only aquatic aspects, here, biota and, specifically, invertebrates. In addition, a separate ecological definition recognizing terrestrial inhabitants has been developed for the alluvial mesovoid shallow substratum beneath dryland non-perennial streams, which flow for a few days each year (Ortuño et al. 2013). Such terrestrial organisms can make significant contributions to biodiversity deep within riverine sediments (Langhans and Tockner 2014). These independent

aquatic and terrestrial conceptualizations are in contrast with Orghidan's (1959, 2010) original recognition that hyporheic conditions transition from saturated (aquatic) to unsaturated (terrestrial) states at ecotone boundaries in perennial streams—and in nonperennial systems, spatial and temporal transitions between wet and dry states greatly increase the extent of ecotonal habitats (Stubbington et al. 2017).

During flowing phases in both perennial and nonperennial streams, upwelling and downwelling zones can support distinct aquatic communities (Datry et al. 2007). Where DO concentrations are relatively high and trophic resources (e.g., particulate and dissolved organic matter) are relatively abundant, such as in downwelling water, many organisms are generalists that primarily inhabit the surface stream and its benthic sediments (Williams and Hynes 1974). Predominantly benthic species include some that use the HZ as a nursery that protects juveniles from stressors, including predation and displacement by flowing water (Giberson and Hall 1988, Feral et al. 2005). In contrast, lower availability of oxygen and trophic resources, as typically characterizes upwelling groundwater, can reduce densities of generalist predators and, thus, enable the persistence of communities dominated by hyporheic and groundwater specialists (Datry et al. 2007), including amphipods, isopods, and a diverse meiofauna (Boulton 2000, Hakenkamp and Palmer 2000). Community composition can also change with depth below the sediment surface, with a shift away from the dominance

of benthic taxa as the influence of surface water and, thus, habitat suitability for these organisms decreases.

When nonperennial streamflow ceases, the shift from lotic to lentic conditions can cause pronounced changes in benthic communities (Bonada et al. 2006, Hill and Milner 2018, Buffagni 2021). Hydrological changes are subdued in the subsurface, where flowing-phase velocities are typically much lower than in the surface stream, and sediment moisture content may remain at or near saturation (Brunke and Gonser 1997; Fig. 5). Hyporheic communities in nonperennial streams have mainly been characterized during flowing phases (Wood et al. 2010, Datry 2012), and biotic responses to subsurface flow cessation have yet to be characterized. As surface water levels decline, mobile organisms that remain in the stream become concentrated within shrinking submerged habitat areas, triggering vertical migrations into the subsurface by organisms seeking refuge from intensifying biotic interactions and causing increased hyporheic densities of primarily benthic organisms (Stubbington et al. 2011, Pařil et al. 2019). If surface water is lost, hyporheic richness can be further enhanced by an influx of benthic taxa seeking refuge from desiccation (Stanley et al. 1994, Clinton et al. 1996). As interstices transition to unsaturated conditions, such taxonomic gains are offset by the loss of desiccation-sensitive organisms. However, whereas benthic community richness nearly always declines with intermittence (Datry et al. 2014, Soria et al. 2017), the number of taxa present in nonperennial HZs may remain stable




Figure 5. Biotic responses to temporal variability in hydrological conditions in the surface stream and benthic and subsurface sediments of nonperennial streams during a typical annual cycle. In a connected hyporheic zone, sediments can experience flowing phases, remain saturated during surface ponded and dry phases, gradually change from saturated to unsaturated then dry conditions after surface drying, then quickly rewet before surface flow resumes. Variability is less pronounced in the subsurface sediments compared with the surface stream and benthic sediments because flow velocities are slower during flowing phases, and interstices remain more humid after free water is lost. Benthic, groundwater, and terrestrial organisms respond to changing conditions differently. Variation in symbol size is proportional to expected densities of the represented organism.

across a gradient of drying duration (Stubbington et al. 2019).

As the benthic zone dries, hyporheic sediments that remain saturated can provide a refuge for aquatic invertebrates, including early instar insects, crustaceans and meiofauna (Clinton et al. 1996, Vander Vorste et al. 2016), as explored particularly through testing of the hyporheic refuge hypothesis (Palmer et al. 1992, Stubbington 2012). Vertebrates, including juvenile lamprey (Rodríguez-Lozano et al. 2019), salamanders (Feral et al. 2005), and small adult fish (Stegman and Minckley 1959, Kawanishi et al. 2013), can also persist in saturated interstices. However, when water levels continue to fall, and where the diminishing dimensions of interstitial pathways prevent vertical migrants from accessing deeper, saturated subsurface sediments, organisms become stranded in humid or dry interstices. Here, sediment characteristics, such as organic matter content, and external influences, including shading and rainfall inputs, can maintain interstitial humidity, allowing subsurface interstices to remain a refuge for aquatic invertebrate and vertebrate life stages with some degree of desiccation tolerance. The abundance and richness of these active or dormant forms relates positively to the moisture content within hyporheic interstices (Stubbington et al. 2009, Datry 2012, Stubbington and Datry 2013).

Desiccation-tolerant aquatic inhabitants of humid hyporheic sediments include both resistant generalists and nonperennial stream specialists. For example, amphipods, including specialist stygobites, can persist for weeks in humid interstices (Gilbert et al. 2018). In addition, Jacobi and Cary (1996) recorded dormant juveniles of 10 stonefly species in the unsaturated subsurface sediments of seasonally nonperennial headwater streams, and Bogan (2017) inferred the dry-phase persistence of the specialist stonefly *Mesocapnia arizonensis* in arid streams with flowing phases as short as 3 mo. Among non-insects, some adult crayfish (DiStefano et al. 2009), frogs (Jared et al. 2020), salamanders, and fish (Secor and Lignot 2010) burrow into deeper, more humid sediments during seasonal dry phases, persisting in either active or dormant states (Secor and Lignot 2010). In addition, annual killifish routinely survive dry phases as dormant eggs within the sediments of ephemeral arid-zone waterbodies (Furness 2016).

As species-specific thresholds are surpassed for core abiotic influences on survival (e.g., water, DO, trophic resources, and temperature), drying sediments can change from a refuge to a graveyard for aquatic organisms (Pařil et al. 2019) and, simultaneously, become a new habitat available for colonization by terrestrial biota. However, whereas recent research has recognized the terrestrial invertebrate biodiversity on dry streambeds (Corti and Datry 2016, Stubbington et al. 2019, Bunting et al. 2021) and within occasionally inundated dryland streambeds (Ortuño et al. 2013), very few studies have explored the terrestrial

communities that move into subsurface sediments in response to declining water levels. Such studies report aquatic–terrestrial assemblages dominated by taxa ranging from fully aquatic (e.g., mites, fly larvae; Bartoszek 2001), semi-aquatic (springtails; Langhans and Tockner 2014), and terrestrial taxa that tolerate inundation (rove beetles; Dieterich 1996).

We also know little about how hyporheic communities respond to flow resumption, with most research considering upward migration of benthic organisms from the subsurface back to the surface stream (Brooks and Boulton 1991, Vander Vorste et al. 2016) rather than characterizing how hyporheic communities colonize and assemble. Assembly may follow similar trajectories to those documented after other disturbances, with faster colonization by benthic compared with groundwater taxa reflecting adaptations that facilitate dispersal in highly dynamic environments (Hancock 2006). In addition, stygobites can colonize quickly where upwelling water passively transports them into groundwater-fed streambeds (Stubbington et al. 2009). Further research is needed to better understand the vertical extent of biodiversity within nonperennial streams, encompassing organisms across the full breadth of environmental preferences from aquatic to terrestrial.

HOW DOES EXISTING NONPERENNIAL STREAM RESEARCH CONSIDER THE HZ?

Above, we present evidence that flowing, ponded, and dry states in nonperennial HZs each have unique, linked hydrological, biogeochemical, and ecological characteristics that interact to contribute to ecosystem processes. This evidence demonstrates that our understanding of these processes—and a broader ecosystem-scale understanding of spatial and temporal variability in HZ structure and function—has been limited by discipline-specific understanding of the HZ, constraining physical and conceptual research. To comprehensively document both disciplinary and cross-discipline biases and limitations, we compiled and examined literature published between 1959 (i.e., the year of Orghidan's seminal paper) and January 2020. We searched for studies of any nonperennial stream (intermittent, ephemeral, temporary) describing sampling of biotic communities in the HZ in any nonperennial stream during a period without surface flow (Table S1). We recorded conditions in which samples were collected and any associated environmental data (e.g., temperature, DO).

We identified 43 primary journal articles published between 1966 and 2020 (Table S1), with most covering a localized spatial scale and with 49 and 33% conducted in North America and Europe, respectively (Fig. S1C). Most studies were conducted in a cool, wet temperate climatic zone (46%), followed by Mediterranean (24%) and arid zones (24%). Two research gaps in particular reflected the

disconnect between discipline-specific conceptualizations of the HZ: 1) few studies explicitly included sampling during the unsaturated, dry, or rewetting phases of the HZ; and 2) only 40% examined how physicochemistry (which, as discussed, varies with hydrology) affects biota during saturated and unsaturated phases. Only a single study (Boulton et al. 1992) characterized the complete range of hydrological conditions within the HZ (high to low infiltration, humid to dry sediments), and only 23% sampled throughout the period in which the surface stream was inundated until dry (Fig. S1D, E). For example, 19% of studies sampled only while water was infiltrating into the HZ, whereas 15% sampled while the HZ had low moisture (Fig. S1E). Only 14% of studies reported sediment moisture content, and most samples were small water volumes collected at a depth of 20 to 50 cm (Fig. S1A, B), regardless of the total extent of the HZ, which was rarely reported. Water characteristics and biota were concurrently measured in 40% of studies, limiting inference of how physicochemical changes, influenced by hydrology, drive biotic patterns. These research gaps reflect both difficulties in sampling the biotic and abiotic conditions within the HZ—especially during unsaturated and dry conditions—and, potentially, a lack of consensus on the relevant parameters to address different ecological research questions in nonperennial streams. It is precisely these gaps regarding connections between the hydrology, biogeochemistry, and biology of the full spatial and temporal extent of the HZ that require further study.

DEFINING AND STANDARDIZING INTERDISCIPLINARY NONPERENNIAL HZ RESEARCH

As nonperennial HZs increase in both spatial and temporal extent because of global change, we call for a return to ideas at the heart of the original conceptualization of the HZ (Orghidan 1959), namely its inclusion of conditions from wet to dry and its consequent support of organisms from aquatic to terrestrial. From its 1st use (Orghidan 1959) and throughout the intervening decades, HZ research has been broad enough to pave the way for its formal extension to include the full range of hydrological, ecological, and biogeochemical conditions experienced in nonperennial systems. Accordingly, we propose a more inclusive definition of nonperennial HZs as the porous subsurface sediments of nonperennial streams, which sometimes directly exchange water, energy, and organisms with adjacent ecosystem components, including the surface channel. This broad definition reflects HZ fluctuations between saturated (lotic or lentic) and unsaturated (humid to dry) interstitial conditions that support organisms with aquatic to terrestrial habitat preferences. Our definition does not constrain the HZ within precise upper or lower boundaries. Instead, during dry phases a gradual decrease in interstitial humidity typically charac-

terizes the vertical transition from benthic to hyporheic sediments. We call for interdisciplinary research spanning the breadth of hydrological conditions experienced within nonperennial HZs, encompassing how biogeochemical processes and ecological communities vary within and between saturated, unsaturated, and dry states.

To promote interdisciplinarity, measurement of a few crucial variables could be standardized across nonperennial HZ research to enable inference of how hydrology influences biogeochemical conditions and ecological communities. Fundamentally, interstitial water requires characterization, ideally including quantification of water movement during saturated conditions and of sediment moisture content during unsaturated states. Although hyporheic water sources and flow directions are difficult to measure, La Montagne et al. (2014) suggest measuring the water level in the stream, adjacent bank, and streambed to determine if the stream is connected to groundwater and to estimate hydraulic gradients that indicate the direction of flow. Simple temperature measurements can be used to trace water movement through the streambed (Constantz 2008), although this method is more complicated when considering lateral fluxes (Shanafield et al. 2010, Xie and Batlle-Aguilar 2017). Automated seepage meters (Solomon et al. 2020) and 3-dimensional sensors (Banks et al. 2018) can also enable rapid capture of hyporheic fluxes at multiple locations within a streambed. During periods in which the stream is not flowing, sediment profiles can be collected (or tensiometers installed in an unsaturated streambed) and analyzed in the laboratory to estimate actual moisture content within the streambed. Geophysical methods, such as electrical resistance tomography, in which low frequency currents are transferred between electrodes and compared, can indicate sediment moisture content, clay content, temperature, and salinity (Ulrich et al. 2015). These methods can be used to characterize whether a stream is connected, disconnected, or in transition and can thereby document both spatial and temporal variation. Lastly, modeling approaches facilitate prediction of the effects of varying hydraulic gradient on physicochemical and biological properties of the HZ (Brunner et al. 2017).

Across hydrological states, DO concentrations and temperature represent key determinants of community composition and, thus, priorities for measurement, ideally by logging data from pre-installed in-situ probes at regular spatial and temporal intervals (e.g., Evans and Petts 1997). Where unsaturated conditions prevent the use of standard meters for in-situ DO measurement, pumping hyporheic water from appropriate depths during both lotic and lentic saturated phases could facilitate collection of sufficiently accurate data to enable comparison among sites and times (Stubbington et al. 2016). Furthermore, sampling campaigns documenting these core abiotic variables should aim to represent the spatial and temporal heterogeneity driven by

interactions between sediment types and water sources by sampling across and within areas with contrasting sediment types and directions/strengths of hydrological exchange. Global-scale studies implemented by interdisciplinary research groups using common standardized protocols have recently generated sufficient data to significantly advance understanding of ecosystem functioning in nonperennial surface streams (e.g., von Schiller et al. 2019), and comparable initiatives could extend into the subsurface.

We also suggest that measurements consider the full spatial and temporal extent of the HZ in its hydrogeological context. The HZ includes the sediments directly beneath the stream and can also extend laterally and stretch to confining layers (Fig. 1A–C). The HZ can also be several meters deep, and if only the shallow HZ is sampled, organisms migrating downward to remain in the wettest available conditions could be missed. Furthermore, physicochemical and biogeochemical conditions that vary, for example, with depth, sediment heterogeneity, and position along a flowpath, could go uncharacterized.

Nonperennial streams may dominate global river networks and are increasing in extent in both space and time. Understanding the linkages between their hydrology, biogeochemistry, and biology is, thus, crucial to inform management strategies that support the structure, function, and integrity of these dynamic ecosystems, including their extensive but often overlooked subsurface components. Our relatively advanced understanding of nonperennial HZs during their saturated and, in particular, their flowing phases covers a subset of their total function. Dry HZs are ecologically active ecosystem components that require greater recognition within nonperennial stream research. We call for research that applies the holistic, interdisciplinary conceptualization of nonperennial HZs developed herein to advance our understanding of these sometimes extensive hidden components of dynamic stream ecosystems as they adapt to global change.

ACKNOWLEDGEMENTS

Author contributions: AGD coordinated manuscript contributions and led the writing. RS, MS, and MAZ assisted with section leadership. All authors contributed to the idea development and writing of the manuscript.

This manuscript is a product of the Dry Rivers Research Coordination Network, which was supported by funding from the National Science Foundation (NSF; DEB 1754389). Additional support to AGD was provided by NSF DEB 1830178.

LITERATURE CITED

- Abril, M., I. Muñoz, and M. Menéndez. 2016. Heterogeneity in leaf litter decomposition in a temporary Mediterranean stream during flow fragmentation. *Science of the Total Environment* 553:330–339.
- Acuña, V., T. Datry, J. Marshall, D. Barcelo, C. N. Dahm, A. Ginebreda, G. McGregor, S. Sabater, K. Tockner, and M. A. Palmer. 2014. Why should we care about temporary waterways? *Science* 343:1080–1081.
- Allen, D. C., D. A. Kopp, K. H. Costigan, T. Datry, B. Hugueny, D. S. Turner, G. S. Bodner, and T. J. Flood. 2019. Citizen scientists document long-term streamflow declines in intermittent rivers of the desert southwest, USA. *Freshwater Science* 38:244–256.
- Ann, Y., K. R. Reddy, and J. J. Delfino. 1999. Influence of redox potential on phosphorus solubility in chemically amended wetland organic soils. *Ecological Engineering* 14:169–180.
- Appelo, C. A. J., and D. Postma. 2004. *Geochemistry, groundwater and pollution*. CRC Press, London, United Kingdom.
- Arias-Real, R., I. Muñoz, C. Gutierrez-Cánovas, V. Granados, P. Lopez-Laseras, and M. Menéndez. 2020. Subsurface zones in intermittent streams are hotspots of microbial decomposition during the non-flow period. *Science of the Total Environment* 703:135485.
- Baldwin, D. S., and A. M. Mitchell. 2000. The effects of drying and re-flooding on the sediment and soil nutrient dynamics of lowland river–floodplain systems: A synthesis. *Regulated Rivers: Research & Management* 16:457–467.
- Banks, E. W., M. A. Shanafield, S. Noorduijn, J. McCallum, J. Lewandowski, and O. Batelaan. 2018. Active heat pulse sensing of 3-D-flow fields in streambeds. *Hydrology and Earth System Sciences* 22:1917–1929.
- Bartoszek, J. E. 2001. Comparison of hyporheic organisms in two intermittent streams to assess a local disturbance. *Journal of Freshwater Ecology* 16:575–579.
- Bencala, K., M. N. Gooseff, and B. A. Kimball. 2011. Rethinking hyporheic flow and transient storage to advance understanding of stream-catchment connections. *Water Resources Research* 47:WR010066.
- Bernhardt, E. S., J. R. Blaszczak, C. D. Ficken, M. L. Fork, K. E. Kaiser, and E. C. Seybold. 2017. Control points in ecosystems: Moving beyond the hot spot hot moment concept. *Ecosystems* 20:665–682.
- Birch, H. F., and M. T. Friend. 1956. Humus decomposition in East African soils. *Nature* 178:500–501.
- Boano, F., J. W. Harvey, A. Marion, A. I. Packman, R. Revelli, L. Ridolfi, and A. Wörman. 2014. Hyporheic flow and transport processes: Mechanisms, models, and biogeochemical implications. *Reviews of Geophysics* 52:603–679.
- Bogan, M. T. 2017. Hurry up and wait: Life cycle and distribution of an intermittent stream specialist (*Mesocapnia arizonensis*). *Freshwater Science* 36:805–815.
- Bonada, N., M. Rieradevall, N. Prat, and V. H. Resh. 2006. Benthic macroinvertebrate assemblages and macrohabitat connectivity in Mediterranean-climate streams of northern California. *Journal of the North American Benthological Society* 25:32–43.
- Boulton, A. J. 2000. The subsurface macrofauna. Pages 337–361 in J. B. Jones and P. J. Mulholland (editors). *Streams and groundwaters*. Academic Press, San Diego, California.
- Boulton, A. J., T. Datry, T. Kasahara, M. Mutz, and J. A. Stanford. 2010. Ecology and management of the hyporheic zone: Stream–groundwater interactions of running waters and their floodplains. *Journal of the North American Benthological Society* 29:26–40.
- Boulton, A. J., E. H. Stanley, S. G. Fisher, and P. S. Lake. 1992. Over-summering strategies of macroinvertebrates in intermittent

- streams in Australia and Arizona. Pages 227–237 in R. D. Robarts and M. L. Bothwell (editors). *Aquatic ecosystems in semi-arid regions: Implications for resource management*. National Hydrology Research Institute Symposium 7.
- Brookfield, A. E., A. T. Hansen, P. L. Sullivan, J. A. Czuba, M. F. Kirk, L. Li, M. E. Newcomer, and G. Wilkinson. 2021. Predicting algal blooms: Are we overlooking groundwater? *Science of the Total Environment* 796:144442.
- Brooks, S. S., and A. J. Boulton. 1991. Recolonization dynamics of benthic macroinvertebrates after artificial and natural disturbances in an Australian temporary stream. *Marine and Freshwater Research* 42:295–308.
- Brunke, M., and T. Gonser. 1997. The ecological significance of exchange processes between rivers and groundwater. *Freshwater Biology* 37:1–33.
- Brunke, M., and T. Gonser. 1999. Hyporheic invertebrates: The clinal nature of interstitial communities structured by hydrological exchange and environmental gradients. *Journal of the North American Benthological Society* 18:344–362.
- Brunner, P., R. Therrien, P. Renard, C. T. Simmons, and H.-J. H. Franssen. 2017. Advances in understanding river–groundwater interactions. *Reviews of Geophysics* 55:818–854.
- Buffagni, A. 2021. The lentic and lotic characteristics of habitats determine the distribution of benthic macroinvertebrates in Mediterranean rivers. *Freshwater Biology* 66:13–34.
- Bunting, G., J. England, K. Gething, T. Sykes, J. Webb, and R. Stubbington. 2021. Aquatic and terrestrial invertebrate community responses to drying in chalk streams. *Water and Environment Journal* 35:229–241.
- Burrows, R. M., L. Beesley, M. M. Douglas, B. J. Pusey, and M. J. Kennard. 2020. Water velocity and groundwater upwelling control benthic algal biomass in a sandy tropical river during base flow: Implications for water resource development. *Hydrobiologia* 847:1207–1219.
- Burrows, R. M., H. Rutledge, N. R. Bond, S. M. Eberhard, A. Auhl, M. S. Andersen, D. G. Valdez, and M. J. Kennard. 2017. High rates of organic carbon processing in the hyporheic zone of intermittent streams. *Scientific Reports* 7:13198.
- Busch, M. H., K. H. Costigan, K. M. Fritz, T. Datry, C. A. Krabbenhoft, J. C. Hammond, M. Zimmer, J. D. Olden, R. M. Burrows, W. K. Dodds, K. S. Boersma, M. Shanafield, S. K. Kampf, M. C. Mims, M. T. Bogan, A. S. Ward, M. Perez Rocha, S. Godsey, G. H. Allen, J. R. Blaszczak, C. N. Jones, and D. C. Allen. 2020. What's in a name? Patterns, trends, and suggestions for defining non-perennial rivers and streams. *Water* 12:1980.
- Cardenas, M. B. 2015. Hyporheic zone hydrologic science: A historical account of its emergence and a prospectus. *Water Resources Research* 51:3601–3616.
- Caruso, A., F. Boano, L. Ridolfi, D. L. Chopp, and A. Packman. 2017. Biofilm-induced bioclogging produces sharp interfaces in hyporheic flow, redox conditions, and microbial community structure. *Geophysical Research Letters* 44:4917–4925.
- Casas-Ruiz, J. P., J. Tittel, D. von Schiller, N. Catalán, B. Obrador, L. Gómez-Gener, E. Zwirnmann, S. Sabater, and R. Marcé. 2016. Drought-induced discontinuities in the source and degradation of dissolved organic matter in a Mediterranean river. *Biogeochemistry* 127:125–139.
- Chafiq, M., J. Gibert, and C. Claret. 1999. Interactions among sediments, organic matter, and microbial activity in the hyporheic zone of an intermittent stream. *Canadian Journal of Fisheries and Aquatic Sciences* 56:487–495.
- Claret, C., and A. J. Boulton. 2008. Integrating hydraulic conductivity with biogeochemical gradients and microbial activity along river–groundwater exchange zones in a subtropical stream. *Hydrogeology Journal* 17:151.
- Clinton, S. M., N. B. Grimm, and S. G. Fisher. 1996. Response of a hyporheic invertebrate assemblage to drying disturbance in a desert stream. *Journal of the North American Benthological Society* 15:700–712.
- Colls, M., X. Timoner, C. Font, S. Sabater, and V. Acuña. 2019. Effects of duration, frequency, and severity of the non-flow period on stream biofilm metabolism. *Ecosystems* 22:1393–1405.
- Constantz, J. 2008. Heat as a tracer to determine streambed water exchanges. *Water Resources Research* 44:WR006996.
- Corti, R., and T. Datry. 2016. Terrestrial and aquatic invertebrates in the riverbed of an intermittent river: Parallels and contrasts in community organisation. *Freshwater Biology* 61:1308–1320.
- Costigan, K. H., M. D. Daniels, and W. K. Dodds. 2015. Fundamental spatial and temporal disconnections in the hydrology of an intermittent prairie headwater network. *Journal of Hydrology* 522:305–316.
- Datry, T. 2012. Benthic and hyporheic invertebrate assemblages along a flow intermittence gradient: Effects of duration of dry events. *Freshwater Biology* 57:563–574.
- Datry, T., N. Bonada, and A. J. Boulton. 2017. *Intermittent rivers and ephemeral streams: Ecology and management*. Academic Press, San Diego, California.
- Datry, T., N. Bonada, and J. Heino. 2016. Towards understanding the organisation of metacommunities in highly dynamic ecological systems. *Oikos* 125:149–159.
- Datry, T., A. J. Boulton, N. Bonada, K. Fritz, C. Leigh, E. Sauquet, K. Tockner, B. Hugueny, and C. N. Dahm. 2018. Flow intermittence and ecosystem services in rivers of the Anthropocene. *Journal of Applied Ecology* 55:353–364.
- Datry, T., and S. T. Larned. 2008. River flow controls ecological processes and invertebrate assemblages in subsurface flowpaths of an ephemeral river reach. *Canadian Journal of Fisheries and Aquatic Sciences* 65:1532–1544.
- Datry, T., S. T. Larned, and M. R. Scarsbrook. 2007. Responses of hyporheic invertebrate assemblages to large-scale variation in flow permanence and surface–subsurface exchange. *Freshwater Biology* 52:1452–1462.
- Datry, T., S. T. Larned, and K. Tockner. 2014. Intermittent rivers: A challenge for freshwater ecology. *BioScience* 64:229–235.
- DelVecchia, A. G., J. A. Stanford, and X. Xu. 2016. Ancient and methane-derived carbon subsidizes contemporary food webs. *Nature Communications* 7:13163.
- Dieterich, M. 1996. Methods and preliminary results from a study on the habitat functions of the gravel bar interior in alluvial floodplains. *Verhandlungen der Gesellschaft für Ökologie* 26:363–367.
- DiStefano, R. J., D. D. Magoulick, E. M. Imhoff, and E. R. Larson. 2009. Imperiled crayfishes use hyporheic zone during seasonal drying of an intermittent stream. *Journal of the North American Benthological Society* 28:142–152.
- Duarte, S., J. Mora-Gómez, A. M. Romaní, F. Cássio, and C. Pascoal. 2017. Responses of microbial decomposers to drought in

- streams may depend on the environmental context. *Environmental Microbiology Reports* 9:756–765.
- Evans, E. C., and G. E. Petts. 1997. Hyporheic temperature patterns within riffles. *Hydrological Sciences Journal* 42:199–213.
- Falke, J. A., K. D. Fausch, R. Magelky, A. Aldred, D. S. Durnford, L. K. Riley, and R. Oad. 2011. The role of groundwater pumping and drought in shaping ecological futures for stream fishes in a dryland river basin of the western Great Plains, USA. *Ecohydrology* 4:682–697.
- Fazi, S., E. Vázquez, E. O. Casamayor, S. Amalfitano, and A. Butturini. 2013. Stream hydrological fragmentation drives bacterioplankton community composition. *PLoS ONE* 8:e64109.
- Febria, C. M., P. Beddoes, R. R. Fulthorpe, and D. D. Williams. 2012. Bacterial community dynamics in the hyporheic zone of an intermittent stream. *The ISME Journal* 6:1078–1088.
- Febria, C. M., J. D. Hosen, B. C. Crump, M. A. Palmer, and D. D. Williams. 2015. Microbial responses to changes in flow status in temporary headwater streams: A cross-system comparison. *Frontiers in Microbiology* 6:522.
- Feral, D., M. A. Camann, and H. H. Welsh Jr. 2005. *Dicamptodon tenebrosus* larvae within hyporheic zones of intermittent streams in California. *Herpetological Review* 36(1):26–27.
- Fleckenstein, J. H., R. G. Niswonger, and G. E. Fogg. 2006. River-aquifer interactions, geologic heterogeneity, and low-flow management. *Groundwater* 44:837–852.
- Furness, A. I. 2016. The evolution of an annual life cycle in killifish: Adaptation to ephemeral aquatic environments through embryonic diapause. *Biological Reviews* 91:796–812.
- Giberson, D. J., and R. J. Hall. 1988. Seasonal variation in faunal distribution within the sediments of a Canadian Shield stream, with emphasis on responses to spring floods. *Canadian Journal of Fisheries and Aquatic Sciences* 45:1994–2002.
- Gilbert, H., J. Keany, and D. C. Culver. 2018. Response of shallow subterranean freshwater amphipods to habitat drying. *Subterranean Biology* 28:15–28.
- Gilbert, J., M. Dole-Olivier, P. Marmonier, and P. Vervier. 1990. Surface water–groundwater ecotones. Pages 199–225 in R. J. Naiman and H. Décamps (editors). *The ecology and management of aquatic-terrestrial ecotones*. Volume 4. The Parthenon Publishing Group, Carnforth, England.
- Gionchetta, G., F. Oliva, M. Menéndez, P. L. Laseras, and A. M. Romani. 2019. Key role of streambed moisture and flash storms for microbial resistance and resilience to long-term drought. *Freshwater Biology* 64:306–322.
- Gionchetta, G., F. Oliva, A. M. Romani, and L. Bañeras. 2020. Hydrological variations shape diversity and functional responses of streambed microbes. *Science of the Total Environment* 714:136838.
- Gómez, R., M. I. Arce, J. J. Sánchez, and M. del Mar Sánchez-Montoya. 2012. The effects of drying on sediment nitrogen content in a Mediterranean intermittent stream: A microcosms study. *Hydrobiologia* 679:43–59.
- Gómez-Gener, L., B. Obrador, R. Marcé, V. Acuña, N. Catalán, J. P. Casas-Ruiz, S. Sabater, I. Muñoz, and D. von Schiller. 2016. When water vanishes: Magnitude and regulation of carbon dioxide emissions from dry temporary streams. *Ecosystems* 19:710–723.
- Gooseff, M. N., D. M. McKnight, R. L. Runkel, and B. H. Vaughn. 2003. Determining long time-scale hyporheic zone flow paths in Antarctic streams. *Hydrological Processes* 17:1691–1710.
- Granados, V., C. Gutiérrez-Cánovas, R. Arias-Real, B. Obrador, A. Harjung, and A. Butturini. 2020. The interruption of longitudinal hydrological connectivity causes delayed responses in dissolved organic matter. *Science of the Total Environment* 713:136619.
- Hakenkamp, C. C., and M. A. Palmer. 2000. The ecology of hyporheic meiofauna. Pages 307–336 in J. B. Jones and P. J. Mulholland (editors). *Streams and ground waters*. Academic Press, San Diego, California.
- Hammond, J., M. Zimmer, M. Shanafield, K. Kaiser, S. Godsey, M. Mims, S. Zipper, R. Burrows, S. Kampf, W. Dodds, C. Jones, C. Krabbenhoft, K. Boersma, T. Datry, J. Olden, G. Allen, A. Price, K. Costigan, R. Hale, and D. Allen. 2021. Spatial patterns and drivers of nonperennial flow regimes in the contiguous United States. *Geophysical Research Letters* 48:GL090794.
- Hancock, P. J. 2006. The response of hyporheic invertebrate communities to a large flood in the Hunter River, New South Wales. *Hydrobiologia* 568:255–262.
- Hancock, P. J., A. J. Boulton, and W. F. Humphreys. 2005. Aquifers and hyporheic zones: Towards an ecological understanding of groundwater. *Hydrogeology Journal* 13:98–111.
- Helton, A. M., M. S. Wright, E. S. Bernhardt, G. C. Poole, R. M. Cory, and J. A. Stanford. 2015. Dissolved organic carbon lability increases with water residence time in the alluvial aquifer of a river floodplain ecosystem. *Journal of Geophysical Research: Biogeosciences* 120:693–706.
- Hensley, R. T., L. Kirk, M. Spangler, M. N. Gooseff, and M. J. Cohen. 2019. Flow extremes as spatiotemporal control points on river solute fluxes and metabolism. *Journal of Geophysical Research: Biogeosciences* 124:537–555.
- Hill, M. J., and V. S. Milner. 2018. Ponding in intermittent streams: A refuge for lotic taxa and a habitat for newly colonising taxa? *Science of the Total Environment* 628:1308–1316.
- Hosen, J. D., K. S. Aho, A. P. Appling, E. C. Creech, J. H. Fair, R. O. Hall Jr, E. D. Kyzivat, R. S. Lowenthal, S. Matt, and J. Morrison. 2019. Enhancement of primary production during drought in a temperate watershed is greater in larger rivers than headwater streams. *Limnology and Oceanography* 64:1458–1472.
- Humphries, P., and D. S. Baldwin. 2003. Drought and aquatic ecosystems: An introduction. *Freshwater Biology* 48:1141–1146.
- Jacobi, G. Z., and S. J. Cary. 1996. Winter stoneflies (Plecoptera) in seasonal habitats in New Mexico, USA. *Journal of the North American Benthological Society* 15:690–699.
- Jared, C., P. L. Mailho-Fontana, J. Mendelson, and M. M. Antoniazzi. 2020. Life history of frogs of the Brazilian semi-arid (Caatinga), with emphasis in aestivation. *Acta Zoologica* 101:302–310.
- Kawanishi, R., M. Inoue, R. Dohi, A. Fujii, and Y. Miyake. 2013. The role of the hyporheic zone for a benthic fish in an intermittent river: A refuge, not a graveyard. *Aquatic Sciences* 75:425–431.
- Kehew, A. E. 2001. *Applied chemical hydrogeology*. Prentice Hall, Upper Saddle River, New Jersey.
- Kiel, B. A., and M. B. Cardenas. 2014. Lateral hyporheic exchange throughout the Mississippi River network. *Nature Geoscience* 7:413–417.

- Krause, S., D. M. Hannah, and J. H. Fleckenstein. 2009. Hyporheic hydrology: Interactions at the groundwater–surface water interface. *Hydrological Processes* 23:2103–2107.
- Krause, S., D. M. Hannah, J. H. Fleckenstein, C. M. Heppell, D. Kaeser, R. Pickup, G. Pinay, A. L. Robertson, and P. J. Wood. 2011. Inter-disciplinary perspectives on processes in the hyporheic zone. *Ecohydrology* 4:481–499.
- La Montagne, S., A. R. Taylor, P. G. Cook, R. S. Crosbie, R. Brownbill, R. M. Williams, and P. Brunner. 2014. Field assessment of surface water–groundwater connectivity in a semi-arid river basin (Murray–Darling, Australia). *Hydrological Processes* 28:1561–1572.
- Langhans, S. D., and K. Tockner. 2014. Is the unsaturated sediment a neglected habitat for riparian arthropods? Evidence from a large gravel-bed river. *Global Ecology and Conservation* 2:129–137.
- Leung, P. M., S. K. Bay, D. V. Meier, E. Chiri, D. A. Cowan, O. Gillor, D. Woebken, and C. Greening. 2020. Energetic basis of microbial growth and persistence in desert ecosystems. *MSystems* 5:e00495-19.
- Lewandowski, J., S. Arnon, E. Banks, O. Batelaan, A. Betterle, T. Broecker, C. Coll, J. D. Drummond, J. G. Garcia, J. Galloway, J. Gomez-Velez, R. C. Grabowski, S. P. Herzog, R. Hinkelmann, A. Höhne, J. Hollender, M. A. Horn, A. Jaeger, S. Krause, A. L. Prats, C. Magliozzi, K. Meinikmann, B. B. Mojarrad, B. M. Mueller, I. Peralta-Maraver, A. L. Popp, M. Posselt, A. Putschew, M. Radke, M. Raza, J. Riml, A. Robertson, C. Rutere, J. L. Schaper, M. Schirmer, H. Schulz, M. Shanafield, T. Singh, A. S. Ward, P. Wolke, A. Wörman, and L. Wu. 2019. Is the Hyporheic zone relevant beyond the scientific community? *Water* 11:2230.
- Mach, V., M. B. Blaser, P. Claus, P. P. Chaudhary, and M. Rulik. 2015. Methane production potentials, pathways, and communities of methanogens in vertical sediment profiles of river Sitka. *Frontiers in Microbiology* 6:506.
- McCallum, J. L., and M. Shanafield. 2016. Residence times of stream–groundwater exchanges due to transient stream stage fluctuations. *Water Resources Research* 52:2059–2073.
- Messenger, M. L., B. Lehner, C. Cockburn, N. Lamouroux, H. Pella, T. Snelder, K. Tockner, T. Trautmann, C. Watt, and T. Datry. 2021. Global prevalence of non-perennial rivers and streams. *Nature* 594:391–397.
- Newcomer, M. E., S. S. Hubbard, J. H. Fleckenstein, U. Maier, C. Schmidt, M. Thullner, C. Ulrich, N. Flipo, and Y. Rubin. 2018. Influence of hydrological perturbations and riverbed sediment characteristics on hyporheic zone respiration of CO₂ and N₂. *Journal of Geophysical Research: Biogeosciences* 123:902–922.
- Orghidan, T. 1959. Ein neuer Lebensraum des unterirdischen Wassers: Der hyporheische Biotop. *Archiv für Hydrobiologie* 55:392–414.
- Orghidan, T. 2010. A new habitat of subsurface waters: The hyporheic biotope. *Fundamental and Applied Limnology: Archiv für Hydrobiologie* 176:291–302.
- Ortuño, V. M., J. D. Gilgado, A. Jiménez-Valverde, A. Sendra, G. Pérez-Suárez, and J. J. Herrero-Borgoñón. 2013. The “alluvial mesovoid shallow substratum”, a new subterranean habitat. *PLoS ONE* 8:e76311.
- Palmer, M. A., A. E. Bely, and K. E. Berg. 1992. Response of invertebrates to lotic disturbance: A test of the hyporheic refuge hypothesis. *Oecologia* 89:182–194.
- Pařil, P., C. Leigh, M. Polářek, R. Sarremejane, P. Řezníčková, A. Dostálová, and R. Stubbington. 2019. Short-term streambed drying events alter amphipod population structure in a central European stream. *Fundamental and Applied Limnology* 193: 51–64.
- Peng, J., B. Wang, Y. Song, P. Yuan, and Z. Liu. 2007. Adsorption and release of phosphorus in the surface sediment of a wastewater stabilization pond. *Ecological Engineering* 31:92–97.
- Perkin, J. S., K. B. Gido, J. A. Falke, K. D. Fausch, H. Crockett, E. R. Johnson, and J. Sanderson. 2017. Groundwater declines are linked to changes in Great Plains stream fish assemblages. *Proceedings of the National Academy of Sciences* 114:7373–7378.
- Quichimbo, E. A., M. B. Singer, and M. O. Cuthbert. 2020. Characterising groundwater–surface water interactions in idealised ephemeral stream systems. *Hydrological Processes* 34:3792–3806.
- Rodríguez-Lozano, P., R. A. Leidy, and S. M. Carlson. 2019. Brook lamprey survival in the dry riverbed of an intermittent stream. *Journal of Arid Environments* 166:83–85.
- Runkel, R. L., D. M. McKnight, and H. Rajaram. 2003. Modeling hyporheic zone processes. *Advances in Water Resources* 26: 901–905.
- Sabater, S., X. Timoner, C. Borrego, and V. Acuña. 2016. Stream biofilm responses to flow intermittency: From cells to ecosystems. *Frontiers in Environmental Science* 4:14.
- Saup, C. M., S. R. Bryant, A. R. Nelson, K. D. Harris, A. H. Sawyer, J. N. Christensen, M. M. Tfaily, K. H. Williams, and M. J. Wilkins. 2019. Hyporheic zone microbiome assembly is linked to dynamic water mixing patterns in snowmelt-dominated headwater catchments. *Journal of Geophysical Research: Biogeosciences* 124:3269–3280.
- Schlesinger, W. H., and E. S. Bernhardt. 2013. Wetland ecosystems. Pages 233–274 in W. H. Schlesinger and E. S. Bernhardt (editors). *Biogeochemistry*. 3rd edition. Academic Press, San Diego, California.
- Secor, S. M., and J.-H. Lignot. 2010. Morphological plasticity of vertebrate aestivation. Pages 183–208 in C. A. Navas and J. E. Carvalho (editors). *Aestivation: Molecular and physiological aspects*. Volume 49. Springer, Heidelberg, Germany.
- Shanafield, M., S. A. Bourke, M. A. Zimmer, and K. H. Costigan. 2021. An overview of the hydrology of non-perennial rivers and streams. *Wiley Interdisciplinary Reviews: Water* 8:e1504.
- Shanafield, M., G. Pohl, and R. Susfalk. 2010. Use of heat-based vertical fluxes to approximate total flux in simple channels. *Water Resources Research* 46:WR007956.
- Siebers, A. R., N. E. Pettit, G. Skrzypek, J. B. Fellman, S. Dogramaci, and P. F. Grierson. 2016. Alluvial ground water influences dissolved organic matter biogeochemistry of pools within intermittent dryland streams. *Freshwater Biology* 61:1228–1241.
- Skoulikidis, N. Th., L. Vardakas, Y. Amaxidis, and P. Michalopoulos. 2017. Biogeochemical processes controlling aquatic quality during drying and rewetting events in a Mediterranean non-perennial river reach. *Science of the Total Environment* 575:378–389.
- Solagaistua, L., M. Arroita, I. Aristi, A. Larrañaga, and A. Elosegi. 2016. Changes in discharge affect more surface than subsurface breakdown of organic matter in a mountain stream. *Marine and Freshwater Research* 67:1826–1834.

- Solomon, D. K., E. Humphrey, T. E. Gilmore, D. P. Genereux, and V. Zlotnik. 2020. An automated seepage meter for streams and lakes. *Water Resources Research* 56:WR026983.
- Soria, M., C. Leigh, T. Datry, L. M. Bini, and N. Bonada. 2017. Biodiversity in perennial and intermittent rivers: A meta-analysis. *Oikos* 126:1078–1089.
- Stanford, J. A., and J. V. Ward. 1988. The hyporheic habitat of river ecosystems. *Nature* 335:64–66.
- Stanford, J. A., and J. V. Ward. 1993. An ecosystem perspective of alluvial rivers: Connectivity and the hyporheic corridor. *Journal of the North American Benthological Society* 12:48–60.
- Stanford, J. A., J. Gibert, and D. Danielopol. 1994. *Groundwater ecology*. Academic Press, San Diego, California.
- Stanley, E. H., D. L. Buschman, A. J. Boulton, N. B. Grimm, and S. G. Fisher. 1994. Invertebrate resistance and resilience to intermittency in a desert stream. *American Midland Naturalist* 131:288–300.
- Stanley, E. H., S. G. Fisher, and N. B. Grimm. 1997. Ecosystem expansion and contraction in streams. *BioScience* 47:427–435.
- Stegman, J. L., and W. L. Minckley. 1959. Occurrence of three species of fishes in interstices of gravel in an area of subsurface flow. *Copeia* 1959:341–341.
- Stonedahl, S. H., J. W. Harvey, A. Wörman, M. Salehin, and A. I. Packman. 2010. A multiscale model for integrating hyporheic exchange from ripples to meanders. *Water Resources Research* 46:WR008865.
- Stubbington, R. 2012. The hyporheic zone as an invertebrate refuge: A review of variability in space, time, taxa and behaviour. *Marine and Freshwater Research* 63:293–311.
- Stubbington, R., M. Acreman, V. Acuña, P. J. Boon, A. J. Boulton, J. England, D. Gilvear, T. Sykes, and P. J. Wood. 2020. Ecosystem services of temporary streams differ between wet and dry phases in regions with contrasting climates and economies. *People and Nature* 2:660–677.
- Stubbington, R., and T. Datry. 2013. The macroinvertebrate seed-bank promotes community persistence in temporary rivers across climate zones. *Freshwater Biology* 58:1202–1220.
- Stubbington, R., M.-J. Dole-Olivier, D. M. P. Galassi, J.-P. Hogan, and P. J. Wood. 2016. Characterization of macroinvertebrate communities in the hyporheic zone of river ecosystems reflects the pump-sampling technique used. *PLoS ONE* 11:e0164372.
- Stubbington, R., J. England, P. J. Wood, and C. E. Sefton. 2017. Temporary streams in temperate zones: Recognizing, monitoring and restoring transitional aquatic-terrestrial ecosystems. *Wiley Interdisciplinary Reviews: Water* 4:e1223.
- Stubbington, R., V. S. Milner, and P. J. Wood. 2019. Flow intermittence in river networks: Understanding the ecohydrological diversity of aquatic-terrestrial ecosystems. *Fundamental and Applied Limnology: Archiv für Hydrobiologie* 193:1–19.
- Stubbington, R., P. J. Wood, and A. J. Boulton. 2009. Low flow controls on benthic and hyporheic macroinvertebrate assemblages during supra-seasonal drought. *Hydrological Processes: An International Journal* 23:2252–2263.
- Stubbington, R., P. J. Wood, I. Reid, and J. Gunn. 2011. Benthic and hyporheic invertebrate community responses to seasonal flow recession in a groundwater-dominated stream. *Ecology* 4:500–511.
- Timoner, X., C. M. Borrego, V. Acuña, and S. Sabater. 2014. The dynamics of biofilm bacterial communities is driven by flow wax and wane in a temporary stream. *Limnology and Oceanography* 59:2057–2067.
- Timoner, X., M. Colls, S. M. Salomón, F. Oliva, V. Acuña, and S. Sabater. 2020. Does biofilm origin matter? Biofilm responses to non-flow period in permanent and temporary streams. *Freshwater Biology* 65:514–523.
- Triska, F. J., V. C. Kennedy, R. J. Avanzino, G. W. Zellweger, and K. E. Bencala. 1989. Retention and transport of nutrients in a third-order stream in northwestern California: Hyporheic processes. *Ecology* 70:1893–1905.
- Tzoraki, O., N. P. Nikolaidis, Y. Amaxidis, and N. Th. Skoulikidis. 2007. In-stream biogeochemical processes of a temporary river. *Environmental Science & Technology* 41:1225–1231.
- Ulrich, C., S. S. Hubbard, J. Florsheim, D. Rosenberry, S. Borglin, M. Trotta, and D. Seymour. 2015. Riverbed clogging associated with a California riverbank filtration system: An assessment of mechanisms and monitoring approaches. *Journal of Hydrology* 529:1740–1753.
- Vander Vorste, R., F. Malard, and T. Datry. 2016. Is drift the primary process promoting the resilience of river invertebrate communities? A manipulative field experiment in an intermittent alluvial river. *Freshwater Biology* 61:1276–1292.
- Vázquez, E., A. M. Román, F. Sabater, and A. Butturini. 2007. Effects of the dry-wet hydrological shift on dissolved organic carbon dynamics and fate across stream-riparian interface in a Mediterranean catchment. *Ecosystems* 10:239–251.
- Verdonschot, R. C. M., A. M. van Oosten-Siedlecka, C. J. F. ter Braak, and P. F. M. Verdonschot. 2015. Macroinvertebrate survival during cessation of flow and streambed drying in a lowland stream. *Freshwater Biology* 60:282–296.
- Vervier, P., J. Gibert, P. Marmonier, and M.-J. Dole-Olivier. 1992. A perspective on the permeability of the surface freshwater-groundwater ecotone. *Journal of the North American Benthological Society* 11:93–102.
- Villeneuve, S., P. G. Cook, M. Shanafield, C. Wood, and N. White. 2015. Groundwater recharge via infiltration through an ephemeral riverbed, central Australia. *Journal of Arid Environments* 117:47–58.
- von Schiller, D., V. Acuña, I. Aristi, M. Arroita, A. Basaguren, A. Bellin, L. Boyero, A. Butturini, A. Ginebreda, E. Kalogianni, A. Larrañaga, B. Majone, A. Martínez, S. Monroy, I. Muñoz, M. Paunović, O. Pereda, M. Petrovic, J. Pozo, S. Rodríguez-Mozaz, D. Rivas, S. Sabater, F. Sabater, N. Skoulikidis, L. Solagaistua, L. Vardakas, and A. Elosegi. 2017. River ecosystem processes: A synthesis of approaches, criteria of use and sensitivity to environmental stressors. *Science of the Total Environment* 596–597:465–480.
- von Schiller, D., V. Acuña, D. Graeber, E. Martí, M. Ribot, S. Sabater, X. Timoner, and K. Tockner. 2011. Contraction, fragmentation and expansion dynamics determine nutrient availability in a Mediterranean forest stream. *Aquatic Sciences* 73:485–497.
- von Schiller, D., T. Datry, R. Corti, A. Foulquier, K. Tockner, R. Marcé, G. García-Baquero, I. Odriozola, B. Obrador, A. Elosegi, C. Mendoza-Lera, M. O. Gessner, R. Stubbington, R. Albariño, D. C. Allen, F. Altermatt, M. I. Arce, S. Arnon, D. Banas, A. Banegas-Medina, E. Beller, M. L. Blanchette, J. F. Blanco-Libreros, J. Blessing, I. G. Boéchat, K. S. Boersma, M. T. Bogan, N. Bonada, N. R. Bond, K. Brintrup, A. Bruder,

- R. M. Burrows, T. Cancellario, S. M. Carlson, S. Cauvy-Fraunié, N. Cid, M. Danger, B. de F. Terra, A. Dehedin, A. M. D. Girolamo, R. del Campo, V. Díaz-Villanueva, C. P. Duerdoth, F. Dyer, E. Faye, C. Febria, R. Figueroa, B. Four, S. Gafny, R. Gómez, L. Gómez-Gener, M. A. S. Graça, S. Guareschi, B. Gücker, F. Hoppeler, J. L. Hwan, S. Kubheka, A. Laini, S. D. Langhans, C. Leigh, C. J. Little, S. Lorenz, J. Marshall, E. J. Martín, A. McIntosh, E. I. Meyer, M. Miliša, M. C. Mlambo, M. Moleón, M. Morais, P. Negus, D. Niyogi, A. Papatheodoulou, I. Pardo, P. Pařil, V. Peřić, C. Piscart, M. Polářek, P. Rodríguez-Lozano, R. J. Rolls, M. M. Sánchez-Montoya, A. Savić, O. Shumilova, A. Steward, A. Taleb, A. Uzan, R. V. Vorste, N. Waltham, C. Woelfle-Erskine, D. Zak, C. Zarfl, and A. Zoppini. 2019. Sediment respiration pulses in intermittent rivers and ephemeral streams. *Global Biogeochemical Cycles* 33:1251–1263.
- Ward, A. S. 2016. The evolution and state of interdisciplinary hyporheic research. *Wiley Interdisciplinary Reviews: Water* 3:83–103.
- Ward, A. S., and A. I. Packman. 2019. Advancing our predictive understanding of river corridor exchange. *WIREs Water* 6: e1327.
- Williams, D., and H. B. N. Hynes. 1974. The occurrence of benthos deep in the substratum of a stream. *Freshwater Biology* 4: 233–256.
- Williams, D. D. 1996. Environmental constraints in temporary fresh waters and their consequences for the insect fauna. *Journal of the North American Benthological Society* 15:634–650.
- Winter, T. C. 1999. Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeology Journal* 7:28–45.
- Woessner, W. W. 2017. Hyporheic zones. Pages 129–157 in F. R. Hauer and G. A. Lamberti (editors). *Methods in stream ecology*. Volume 1. 3rd edition. Academic Press, San Diego, California.
- Wondzell, S. M., and F. J. Swanson. 1999. Floods, channel change, and the hyporheic zone. *Water Resources Research* 35:555–567.
- Wood, P. J., A. J. Boulton, S. Little, and R. Stubbington. 2010. Is the hyporheic zone a refugium for aquatic macroinvertebrates during severe low flow conditions? *Fundamental and Applied Limnology: Archiv für Hydrobiologie* 176:377–390.
- Wroblicky, G. J., M. E. Campana, H. M. Valett, and C. N. Dahm. 1998. Seasonal variation in surface–subsurface water exchange and lateral hyporheic area of two stream–aquifer systems. *Water Resources Research* 34:317–328.
- Xie, Y., and J. Battle-Aguilar. 2017. Limits of heat as a tracer to quantify transient lateral river–aquifer exchanges. *Water Resources Research* 53:7740–7755.
- Zeglin, L. H., C. N. Dahm, J. E. Barrett, M. N. Gooseff, S. K. Fitzpatrick, and C. D. Takacs-Vesbach. 2011. Bacterial community structure along moisture gradients in the parafluvial sediments of two ephemeral desert streams. *Microbial Ecology* 61:543–556.
- Zimmer, M. A., and L. K. Lautz. 2014. Temporal and spatial response of hyporheic zone geochemistry to a storm event. *Hydrological Processes* 28:2324–2337.
- Zlatanović, S., J. Fabian, K. Premke, and M. Mutz. 2018. Shading and sediment structure effects on stream metabolism resistance and resilience to infrequent droughts. *The Science of the Total Environment* 621:1233–1242.